arXiv:0902.1383v1 [math.RT] 9 Feb 2009

On covariant functions and distributions under the action of a compact group

Anouar SAIDI [*]

February 9, 2009

Abstract. Let G be a compact subgroup of $GL_n(\mathbb{R})$ acting linearly on a finite dimensional vector space E. B. Malgrange has shown that the space $\mathcal{C}^\infty(\mathbb{R}^n, E)^G$ of $\mathcal{C}^\infty$ and G-covariant functions is a finite module over the ring $\mathcal{C}^\infty(\mathbb{R}^n)^G$ of $\mathcal{C}^\infty$ and G-invariant functions. First, we generalize this result for the Schwartz space $\mathscr{S}(\mathbb{R}^n, E)^G$ of G-covariant functions. Secondly, we prove that any G-covariant distribution can be decomposed into a sum of G-invariant distributions multiplied with a fixed family of G-covariant polynomials. This gives a generalization of an Oksak result proved in ([O]).

Keywords : Covariant functions, covariant distributions, compact group.
Subject classification : 46F05, 58C99, 58C81, 58C46.

1 Introduction

Let G be a compact Lie group acting linearly on $\mathbb{R}^n$ and ρ a representation of G in a finite dimensional complex vector space E.

Let $\mathscr{A}(\mathbb{R}^n, E)$ be either of the spaces $\mathcal{C}^\infty(\mathbb{R}^n, E)$, $\mathcal{C}_c^\infty(\mathbb{R}^n, E)$ or $\mathscr{S}(\mathbb{R}^n, E)$. A function f in $\mathscr{A}(\mathbb{R}^n, E)$ is said to be G-covariant if it verifies

$$\rho(g)^{-1} \circ f \circ g = f, \quad \text{for all } g \in G.$$

We denote by $\mathscr{A}(\mathbb{R}^n, E)^G$ the space of G-covariant functions and by $\mathscr{A}(\mathbb{R}^n)^G$ the space of G-invariant functions.

Let E^* be the dual space of E and ρ^* the adjoint representation of ρ in E^*. We consider $\mathscr{A}'(\mathbb{R}^n, E)$ the dual space of $\mathscr{A}(\mathbb{R}^n, E^*)$. A distribution T in $\mathscr{A}'(\mathbb{R}^n, E)$ is said to be G-covariant if it verifies

$$\langle T, \rho(g)^{-1} \circ f \circ g \rangle = \langle T, f \rangle, \quad \text{for all } f \in \mathscr{A}(\mathbb{R}^n, E^*) \text{ and } g \in G.$$

---

[*] Faculty of Sciences of Monastir. Department of Mathematics. 5019 Monastir.Tunisia.
Email: saidi.anouar@yahoo.fr

Let $\mathscr{A}'(\mathbb{R}^n, E)^G$ be the space of G-covariant distributions and $\mathscr{A}'(\mathbb{R}^n)^G$ the space of G-invariant distributions.

Representations of G-covariant functions and distributions in terms of G-covariant polynomials are of frequent use in theoretical physics. B. Malgrange (see [P] of V. Poénaru) has obtained such representation for functions in $\mathcal{C}^\infty(\mathbb{R}^n, E)^G$. More precisely, he has proved that there exists a family of G-covariant polynomials $P_1, \ldots, P_k$ such that every function f in $\mathcal{C}^\infty(\mathbb{R}^n, E)^G$ can be represented in the form

$$f = f_1 P_1 + \ldots + f_r P_k, \qquad \text{where } f_1, \ldots, f_k \in \mathcal{C}^\infty(\mathbb{R}^n)^G. \tag{1.1}$$

A.I. Oksak in [O] has considered the case when the subspace $\mathscr{P}(\mathbb{R}^n, E)^G$ of G-covariant polynomials is a free module over the ring $\mathscr{P}(\mathbb{R}^n)^G$ of G-invariant polynomials. This is equivalent to say that the map $I : (\mathscr{P}(\mathbb{R}^n)^G)^k \longrightarrow \mathscr{P}(\mathbb{R}^n, E)^G$ defined by

$$(p_1, \ldots, p_k) \longmapsto \sum_{i=1}^{k} p_i \; P_i,$$

is an isomorphism. Under this assumption, he established that the map $\Phi : (\mathscr{A}(\mathbb{R}^n)^G)^k \longrightarrow \mathscr{A}(\mathbb{R}^n, E)^G$ defined by

$$(f_1, \ldots, f_k) \longmapsto \sum_{i=1}^{k} f_i \; P_i,$$

is a topological isomorphism when the vector space $\mathscr{A}(\mathbb{R}^n, E)^G$ is equal to $\mathcal{C}^\infty(\mathbb{R}^n, E)^G$, $\mathcal{C}_c^\infty(\mathbb{R}^n, E)^G$ or $\mathscr{S}(\mathbb{R}^n, E)^G$.

A.I. Oksak has also studied the representation of G-covariant distributions. He has proved, under the above assumption, that every distribution T in $\mathscr{A}'(\mathbb{R}^n, E)^G$ can be represented in the form

$$T = \sum_{i=1}^{k} T_i \; P_i, \qquad \text{where } T_1, \ldots, T_k \in \mathscr{A}'(\mathbb{R}^n)^G. \tag{1.2}$$

Our aim in this paper is to generalize the results of B. Malgrange and A.I. Oksak. In section 2, we prove the analogue of the result of B. Malgrange in the spaces $\mathcal{C}_c^\infty(\mathbb{R}^n, E)^G$ and $\mathscr{S}(\mathbb{R}^n, E)^G$. In section 3, we establish the decomposition (1.2) without any assumption on the action of the compact group G.

2 Representation of G-covariant functions

Let G be a compact Lie group and μ its normalized Haar measure. We suppose that G acts linearly on $\mathbb{R}^n$ and we fix an euclidean G-invariant norm on $\mathbb{R}^n$. Let ρ be a representation of G on a finite dimensional complex vector space E. We choose a G-invariant norm $|| \ \ ||_E$ on E. By duality, G acts on E^* the dual space of E. We consider $\mathcal{C}_c^\infty(\mathbb{R}^n, E)$ the space of compact support functions in $\mathcal{C}^\infty(\mathbb{R}^n, E)$. For a compact K in $\mathbb{R}^n$, $\mathcal{C}_K^\infty(\mathbb{R}^n, E)$ denotes the space of functions in $\mathcal{C}_c^\infty(\mathbb{R}^n, E)$ such that $\mathrm{supp}(f) \subset K$ and $\mathscr{S}(\mathbb{R}^n, E)$ the space of functions f in $\mathcal{C}^\infty(\mathbb{R}^n, E)$ satisfying

$$\sup_{x \in \mathbb{R}^n} (1 + ||x||^2)^m \ ||D^\alpha f(x)||_E < \infty,$$

for all $m \in \mathbb{N}$ and $\alpha \in \mathbb{N}^n$.

We equip the spaces $\mathcal{C}^\infty(\mathbb{R}^n, E)$, $\mathcal{C}_c^\infty(\mathbb{R}^n, E)$ and $\mathscr{S}(\mathbb{R}^n, E)$ with their usual topologies. For $g \in G$ and $f \in \mathscr{A}(\mathbb{R}^n, E)$, we denote by $g \cdot f$ the function $\rho(g)^{-1} \circ f \circ g$ which also belongs to $\mathscr{A}(\mathbb{R}^n, E)$. We say that a function f in $\mathscr{A}(\mathbb{R}^n, E)$ is G-covariant if it verifies

$$g \cdot f = f, \quad \text{for all } g \in G.$$

It is clear that $\mathscr{A}(\mathbb{R}^n, E)^G$ is an $\mathscr{A}(\mathbb{R}^n)^G$-module. Our aim in this section is to prove that this module is generated by a finite family of G-covariant polynomials. The case $\mathcal{C}^\infty(\mathbb{R}^n, E)$, is already established by B. Malgrange (see [P]) and the same proof is valid to establish that the set of G-covariant polynomials is a module of finite type over the ring of G-invariant polynomials. We fix $P_1, \ldots, P_k$ a family of G-covariant homogeneous polynomials which generate this module. So we have :

Lemma 2.1 *For all $f \in \mathcal{C}^\infty(\mathbb{R}^n, E)^G$ there exists a family of functions $f_1, \ldots, f_k$ in $\mathcal{C}^\infty(\mathbb{R}^n)^G$ such that*

$$f = \sum_{i=1}^k f_i \ P_i.$$

As corollary of the above lemma, we have

Corollary 2.2 *For all $f \in \mathcal{C}_c^\infty(\mathbb{R}^n, E)^G$ there exists a family of functions $f_1, \ldots, f_k$ in $\mathcal{C}_c^\infty(\mathbb{R}^n)^G$ such that*

$$f = \sum_{i=1}^k f_i \ P_i.$$

The main result of this section is :

Theorem 2.3 *For all* $f \in \mathscr{S}(\mathbb{R}^n, E)^G$ *there exists a family of functions* $f_1, \ldots, f_k$ *in* $\mathscr{S}(\mathbb{R}^n)^G$ *such that*

$$f = \sum_{i=1}^{k} f_i \; P_i.$$

To prove this theorem, we need the following lemmas.

For all $0 \leq r < R$, we consider the sets $B(0,r) = \{x \in \mathbb{R}^n, \; ||x|| < r\}$, $S(0,r) = \{x \in \mathbb{R}^n, \; ||x|| = r\}$, $\overline{B}(0,r) = B(0,r) \cup S(0,r)$ and $C(r,R) = \{x \in \mathbb{R}^n, \; r \leq ||x|| \leq R\}$.

As an improvement of Corollary 2.2, we have

Lemma 2.4 *Let* f *be in* $\mathcal{C}^{\infty}_{\overline{B}(0,1)}(\mathbb{R}^n, E)^G$. *There exists a family of functions* $f_1, \ldots, f_k$ *in* $\mathcal{C}^{\infty}_{\overline{B}(0,1)}(\mathbb{R}^n)^G$ *such that*

$$f = \sum_{i=1}^{k} f_i \; P_i.$$

Proof

Let f be in $\mathcal{C}^{\infty}_{\overline{B}(0,1)}(\mathbb{R}^n, E)^G$. By Corollary 2.2 we have

$$f = \sum_{i=1}^{k} f_i \; P_i, \qquad \text{with } f_1, \ldots, f_k \in \mathcal{C}^{\infty}_c(\mathbb{R}^n)^G.$$

We have to prove that the functions $f_1, \ldots, f_k$ can be chosen with supports in $\overline{B}(0,1)$. For this, we need to use Lemma 8.1.1 of [B]. This lemma treats the case $E = \mathbb{C}$ but it is also true for any finite dimensional vector space.

Let γ be in $\mathcal{C}^{\infty}(\mathbb{R}^n)^G$ satisfying $\gamma \equiv 1$ on $\overline{B}(0, \frac{1}{2})$ and $\text{supp}(\gamma) \subset \overline{B}(0, \frac{2}{3})$. Then $f = \gamma f + (1-\gamma) f$.

The function $(1-\gamma)\; f \in \mathcal{C}^{\infty}_{C(\frac{1}{2},1)}(\mathbb{R}^n, E)^G$, so Lemma 8.1.1 of [B] implies that

$$(1-\gamma(x)) \; f(x) = \sum_{j=1}^{r} g_j(||x||^2) \; \varphi_j(x), \qquad \text{for all } x \in \mathbb{R}^n,$$

with $g_j \in \mathcal{C}^{\infty}_{[\frac{1}{4},1]}(\mathbb{R})$ and $\varphi_j \in \mathcal{C}^{\infty}_{C(\frac{1}{2},1)}(\mathbb{R}^n, E)$. Since G is a compact group, we can suppose that the functions φ_j are G-covariant. Then by Corollary 2.2, we have

$$\varphi_j = \sum_{i=1}^{k} \varphi_{ij} \; P_i, \qquad \text{for all } 1 \leq j \leq r,$$

where $\varphi_{ij} \in \mathcal{C}_c^\infty(\mathbb{R}^n)^G$. So we get

$$(1-\gamma(x))\ f(x) = \sum_{i=1}^{k} \left(\sum_{j=1}^{r} g_j(||x||^2)\varphi_{ij}(x) \right) P_i(x), \quad \text{for all } x \in \mathbb{R}^n.$$

The result follows for the function $(1-\gamma)f$ since the functions $g_j(||\ .\ ||^2)\ \varphi_{ij}$ are in $\mathcal{C}^\infty_{\overline{B}(0,1)}(\mathbb{R}^n)^G$.

According to Corollary 2.2, there exist $h_1, \dots, h_k \in \mathcal{C}_c^\infty(\mathbb{R}^n)^G$ such that

$$\gamma f = \sum_{i=1}^{k} h_i\ P_i.$$

Let θ be in $\mathcal{C}_c^\infty(\mathbb{R}^n)^G$ satisfying $\theta \equiv 1$ on $\overline{B}(0, \frac{2}{3})$ and $\text{supp}(\theta) \subset \overline{B}(0,1)$. It is clear that

$$\gamma f = \theta\ \gamma\ f = \sum_{i=1}^{k} (\theta\ h_i)\ P_i,$$

with $\theta\ h_i \in \mathcal{C}^\infty_{\overline{B}(0,1)}(\mathbb{R}^n)^G$. This completes the proof. $\square$

Lemma 2.5 *Let h be in $\mathcal{C}^\infty(\mathbb{R}^n, E)^G$ such that for all α in $\mathbb{N}^n$, $D^\alpha h \equiv 0$ on $S(0,1)$. Then, for all p in $\mathbb{N}$,*

$$\lim_{||y|| \to 1^-} \frac{h(y)}{(1-||y||^2)^p} = 0.$$

Proof

Let h be in $\mathcal{C}^\infty(\mathbb{R}^n, E)^G$ and p in $\mathbb{N}$. We set $M = \sup\limits_{z \in \overline{B}(0,2)} ||D^{p+1}h(z)||$. By Taylor formula we have

$$h(y) = \int_0^1 \frac{(1-t)^p}{p!}\ D^{p+1}h(y_0 + t(y-y_0)) \cdot (y-y_0)^{(p+1)}\ dt,$$

for all $y \in \overline{B}(0,1)$ and y_0 in $S(0,1)$. Then,

$$|h(y)| \leq M\ ||y-y_0||^{p+1}, \quad \text{for all } y \in \overline{B}(0,1) \text{ and } y_0 \in S(0,1).$$

Let $y \in B(0,1) \setminus \{0\}$ and $y_0 = \frac{y}{\|y\|}$. Then $\|y-y_0\| = 1 - \|y\|$ and

$$\frac{|h(y)|}{(1-||y||^2)^p} = \frac{|h(y)|}{(1+||y||)^p ||y-y_0||^p} \leq M\ ||y-y_0||.$$

This gives the result since

$$\lim_{||y||\to 1^-} ||y-y_0|| = 0.$$

$\square$

The function $\varphi : x \longmapsto \frac{x}{\sqrt{1-\|x\|^2}}$ is a bijection from $B(0,1)$ to $\mathbb{R}^n$ and its inverse function is

$$\psi : x \longmapsto \frac{x}{\sqrt{1+||x||^2}}.$$

For f in $\mathscr{S}(\mathbb{R}^n, E)$, we denote by $I(f)$ the function on $\mathbb{R}^n$ defined by :

$$I(f)(x) = \begin{cases} f\circ\varphi(x) & \text{if } x \in B(0,1) \\ 0 & \text{otherwise} \end{cases} .$$

The map I is one to one and satisfies :

Lemma 2.6 *For all $f \in \mathscr{S}(\mathbb{R}^n, E)$, the function $I(f)$ is in $\mathcal{C}^\infty_{\overline{B}(0,1)}(\mathbb{R}^n, E)$ and the map $I : \mathscr{S}(\mathbb{R}^n, E) \longrightarrow \mathcal{C}^\infty_{\overline{B}(0,1)}(\mathbb{R}^n, E)$ is a topological isomorphism.*

Proof
Let f be in $\mathscr{S}(\mathbb{R}^n, E)$. To prove that $I(f)$ is $\mathcal{C}^\infty$ it is enough to show that

$$\lim_{\|x\|\to 1^-} D^\alpha(f\circ\varphi)(x) = 0, \qquad \text{for all } \alpha \in \mathbb{N}^n.$$

It is clear that, for all $x \in B(0,1)$, $D^\alpha(f\circ\varphi)(x)$ can be written as a sum of elements of the form

$$D^\gamma f(\varphi(x))\; D^{\lambda_1}\varphi_{i_1}(x)\cdots D^{\lambda_r}\varphi_{i_r}(x),$$

with $\lambda_i, \gamma \in \mathbb{N}^n$, and

$$D^{\lambda_j}\varphi_{i_j}(x) = \frac{Q_{i,j}(x)}{(1-||x||^2)^s},$$

where $Q_{i,j}$ is a polynomial and $s \in \frac{1}{2}\mathbb{N}$. Since

$$1-||x||^2 = \frac{1}{1+||\varphi(x)||^2},$$

then there exist a polynomial P and $s' \in \frac{1}{2}\mathbb{N}$ such that

$$D^\gamma f(\varphi(x))\; D^{\lambda_1}\varphi_{i_1}(x)\cdots D^{\lambda_r}\varphi_{i_r}(x) = P(x)\;(1+||\varphi(x)||^2)^{s'}\; D^\gamma f(\varphi(x)). \tag{2.1}$$

As $f \in \mathscr{S}(\mathbb{R}^n, E)$ and $\lim_{||x|| \to 1^-} ||\varphi(x)|| = +\infty$, then

$$\lim_{||x|| \to 1^-} D^\alpha (f \circ \varphi)(x) = 0,$$

and we deduce that $I(f) \in \mathcal{C}^\infty_{\overline{B}(0,1)}(\mathbb{R}^n, E)$.

To establish that I is onto we will prove that if $g \in \mathcal{C}^\infty_{\overline{B}(0,1)}(\mathbb{R}^n, E)$, then $g \circ \psi \in \mathscr{S}(\mathbb{R}^n, E)$ and $I(g \circ \psi) = g$. We fix $r \in \mathbb{N}$ and $\beta \in \mathbb{N}^n$. As in (2.1) there exist $m \in \mathbb{Z}$ and $c > 0$ such that

$$(1 + ||x||^2)^r ||D^\beta (g \circ \psi)(x)||_E \leq c\, (1 + ||x||^2)^m \sup_{\gamma \leq \beta} ||D^\gamma g(\psi(x))||_E.$$

Since $\lim\limits_{||x|| \to +\infty} ||\psi(x)|| = 1$ and $1 + ||x||^2 = \dfrac{1}{1 - ||\psi(x)||^2}$, then Lemma 2.5 insures that

$$\lim_{||x|| \to +\infty} (1 + ||x||^2)^m ||D^\gamma g(\psi(x))||_E = 0,$$

and we deduce that

$$\lim_{||x|| \to +\infty} (1 + ||x||^2)^r ||D^\beta (g \circ \psi)(x))||_E = 0.$$

Hence $g \circ \psi \in \mathscr{S}(\mathbb{R}^n, E)$, and it is easy to verify that $I(g \circ \psi) = g$.

It follows from the closed graph theorem that I is a topological isomorphism. ☐

Proof of Theorem 2.3

Let f be in $\mathscr{S}(\mathbb{R}^n, E)^G$. Since we have

$$\varphi(g\, x) = g.(\varphi\, (x)), \quad \text{for all } g \in G \text{ and } x \in B(0,1),$$

then the function $I(f)$ is in $\mathcal{C}^\infty_{\overline{B}(0,1)}(\mathbb{R}^n, E)^G$ and by Lemma 2.4, we have

$$I(f)(x) = \sum_{i=1}^k g_i(x)\, P_i(x), \quad \text{for all } x \in \mathbb{R}^n,$$

with $g_1, \ldots, g_k \in \mathcal{C}^\infty_{\overline{B}(0,1)}(\mathbb{R}^n)^G$. So for all y in $\mathbb{R}^n$, we have

$$f(y) = I(f)(\psi(y)) = \sum_{i=1}^k g_i(\psi(y))\, P_i(\psi(y)).$$

Let $s_1, \ldots, s_k$ be the degrees of homogeneity of $P_1, \ldots, P_k$. Then

$$P_i(\psi(y)) = P_i\big(\frac{y}{\sqrt{1 + ||y||^2}}\big) = \big(\frac{1}{\sqrt{1 + ||y||^2}}\big)^{s_i} P_i(y), \quad \text{for all } 1 \leq i \leq k.$$

Hence

$$f(y) = \sum_{i=1}^{k} f_i(y)\ P_i(y),$$

with

$$f_i(y) = \Big(\frac{1}{\sqrt{1+||y||^2}}\Big)^{s_i}\ g_i\Big(\frac{y}{\sqrt{1+||y||^2}}\Big) \in \mathscr{S}(\mathbb{R}^n)^G.$$

To conclude, we have to prove that $f_1, \ldots, f_k \in \mathscr{S}(\mathbb{R}^n)$. We fix $1 \leq i \leq k$ and we consider the function $h_i : \mathbb{R}^n \longrightarrow \mathbb{C}$ defined by

$$h_i(x) = \begin{cases} (\sqrt{1-||x||^2})^{s_i}\ g_i(x) & \text{if } x \in B(0,1) \\ 0 & \text{otherwise} \end{cases}.$$

The partial derivatives of h_i in $B(0,1)$ are of the form $P(x)(1-||x||^2)^r D^\gamma g_i(x)$ with $r \in \frac{1}{2}\mathbb{Z}$, $\gamma \in \mathbb{N}^n$ and P a polynomial. Hence, by Lemma 2.5, we conclude that $\lim\limits_{||x|| \to 1^-} D^\alpha h_i(x) = 0$ for all $\alpha \in \mathbb{N}^n$ and so $h_i \in \mathcal{C}^\infty_{\overline{B}(0,1)}(\mathbb{R}^n)$. Finally, we verify that $f_i = h_i \circ \psi = I^{-1}(h_i)$ and that f_i is G-invariant. This completes the proof. $\square$

3 Representation of G-covariant distributions

Let $\mathscr{A}'(\mathbb{R}^n, E)$ be the topological dual of the space $\mathscr{A}(\mathbb{R}^n, E^*)$ equipped with its weak topology. A distribution T in $\mathscr{A}'(\mathbb{R}^n, E)$ is said to be G-covariant if it verifies

$$\langle T, g \cdot f \rangle = \langle T, f \rangle, \quad \text{for all } f \in \mathscr{A}(\mathbb{R}^n, E^*) \text{ and } g \in G. \tag{3.1}$$

We denote by $\mathscr{A}'(\mathbb{R}^n, E)^G$ the space of G-covariant distributions. In the particular case of the trivial representation, a G-covariant distribution T is G-invariant and we denote by $\mathscr{A}'(\mathbb{R}^n)^G$ the space $\mathscr{A}'(\mathbb{R}^n, \mathbb{C})^G$.

For $f \in \mathscr{A}(\mathbb{R}^n, E^*)$ and $h \in \mathscr{A}(\mathbb{R}^n, E)$, we define the function $\langle f, h \rangle$ in $\mathscr{A}(\mathbb{R}^n)$ by

$$\langle f, h \rangle(x) = \langle f(x), h(x) \rangle, \quad \text{for all } x \in \mathbb{R}^n.$$

Let $\theta \in \mathscr{A}'(\mathbb{R}^n)$ and $h \in \mathscr{A}(\mathbb{R}^n, E)$. We denote by $\theta\, h$ the distribution in $\mathscr{A}'(\mathbb{R}^n, E)$ defined by

$$\langle \theta\ h, f \rangle = \langle \theta, \langle f, h \rangle \rangle, \quad \text{for all } f \in \mathscr{A}(\mathbb{R}^n, E^*).$$

If $\theta \in \mathscr{A}'(\mathbb{R}^n)^G$ and $h \in \mathscr{A}(\mathbb{R}^n, E)^G$, then $\theta\, h \in \mathscr{A}'(\mathbb{R}^n, E)^G$.

Then we have the following theorem :

Theorem 3.1 *For all θ in $\mathscr{D}'(\mathbb{R}^n, E)^G$ there exists a family of distributions $\theta_1, \ldots, \theta_k$ in $\mathscr{D}'(\mathbb{R}^n)^G$ such that $\theta = \sum_{i=1}^{k} \theta_i \, P_i$.*

Before the proof of this theorem, we establish some lemmas.

For $\varphi \in \mathcal{C}_c^\infty(\mathbb{R}^n, E)$, let T_φ be the distribution in $\mathscr{D}'(\mathbb{R}^n, E)$ defined by $T_\varphi(f) = \int_{\mathbb{R}^n} \langle \varphi(x), f(x) \rangle \, dx$. Then we can identify $\mathcal{C}_c^\infty(\mathbb{R}^n, E)$ with a subspace of $\mathscr{D}'(\mathbb{R}^n, E)$. Moreover, if φ is a G-covariant function, then T_φ is a G-covariant distribution and we have the following result.

Lemma 3.2 *The space $\mathcal{C}_c^\infty(\mathbb{R}^n, E)^G$ is dense in $\mathscr{D}'(\mathbb{R}^n, E)^G$.*

Proof
Let f be in $\mathscr{D}(\mathbb{R}^n, E^*)$. The map $g \longmapsto g \cdot f$ from G to $\mathscr{D}(\mathbb{R}^n, E^*)$ is clearly continuous. By Proposition 5 in [BO], there exists a unique function $\mathscr{M}(f)$ in $\mathscr{D}(\mathbb{R}^n, E^*)$, such that for all $T \in \mathscr{D}'(\mathbb{R}^n, E)$, we have

$$\langle T, \mathscr{M}(f) \rangle = \int_G \langle T, g \cdot f \rangle \, d\mu(g). \tag{3.2}$$

It follows from this equality that

$$\mathscr{M}(f) = \mathscr{M}(g \cdot f), \quad \text{for all } f \in \mathscr{D}(\mathbb{R}^n, E^*) \text{ and all } g \in G. \tag{3.3}$$

For $T = \delta_x$, the Dirac distribution on $x \in \mathbb{R}^n$, the formula (3.2) gives

$$\mathscr{M}(f)(x) = \int_G g \cdot f \; (x) \; d\mu(g).$$

Then we can verify that :

i) $\mathscr{M}(f) \in \mathscr{D}(\mathbb{R}^n, E^*)^G$.

ii) $\mathscr{M}(f) = f$ if f is G-covariant.

iii) The linear map $\mathscr{M} : \mathscr{D}(\mathbb{R}^n, E^*) \longrightarrow \mathscr{D}(\mathbb{R}^n, E^*)$ is continuous (according to the graph theorem).

iv) $\mathscr{M}(\mathcal{C}_c^\infty(\mathbb{R}^n, E)) = \mathcal{C}_c^\infty(\mathbb{R}^n, E)^G$.

Let ${}^t\mathscr{M}$ be the transpose map of $\mathscr{M}$. It is continuous for the weak topologies and verifies

$${}^t\mathscr{M}(T) = T, \quad \text{for all } T \in \mathscr{D}'(\mathbb{R}^n, E)^G, \tag{3.4}$$

and

$$ {}^t\mathscr{M}(T_\varphi) = T_{\mathscr{M}(\varphi)}, \qquad \text{for all } \varphi \in \mathcal{C}_c^\infty(\mathbb{R}^n, E). \tag{3.5} $$

Since $\mathcal{C}_c^\infty(\mathbb{R}^n, E)$ is dense in $\mathscr{A}'(\mathbb{R}^n, E)$, then ${}^t\mathscr{M}(\mathcal{C}_c^\infty(\mathbb{R}^n, E))$ is dense in ${}^t\mathscr{M}(\mathscr{A}'(\mathbb{R}^n, E))$.

To establish the lemma we will prove that ${}^t\mathscr{M}(\mathscr{A}'(\mathbb{R}^n, E)) = \mathscr{A}'(\mathbb{R}^n, E)^G$ and ${}^t\mathscr{M}(\mathcal{C}_c^\infty(\mathbb{R}^n, E)) = \mathcal{C}_c^\infty(\mathbb{R}^n, E)^G$.

The inclusion ${}^t\mathscr{M}(\mathscr{A}'(\mathbb{R}^n, E)) \subset \mathscr{A}'(\mathbb{R}^n, E)^G$ follows from the formula (3.3) and the formula (3.4) gives the inverse inclusion. Finally, the equality ${}^t\mathscr{M}(\mathcal{C}_c^\infty(\mathbb{R}^n, E)) = \mathcal{C}_c^\infty(\mathbb{R}^n, E)^G$ follows from the formula (3.5). □

Let $P_{\mathscr{A}} : \mathscr{A}(\mathbb{R}^n, E^*) \longrightarrow (\mathscr{A}(\mathbb{R}^n))^k$ be the continuous linear map defined by

$$ P_{\mathscr{A}}(f) = (\langle f, P_1 \rangle, \ldots, \langle f, P_k \rangle), $$

and ${}^tP_{\mathscr{A}}$ its transpose. Let $(e_i)_{1 \le i \le d}$ be a basis of E and $(e_i^*)_{1 \le i \le d}$ its dual basis. For $1 \le l \le k$, we have $P_l = \sum_{i=1}^d P_{li}\, e_i$ where $(P_{li})_{1 \le i \le d}$ are homogeneous polynomials with the same degree of homogeneity s_l as P_l.

For all function f in $\mathscr{A}(\mathbb{R}^n, E^*)$, there exist $f_1, \ldots, f_d \in \mathscr{A}(\mathbb{R}^n)$ such that $f = \sum_{i=1}^d f_i\, e_i^*$. Then the spaces $\mathscr{A}(\mathbb{R}^n, E^*)$ and $(\mathscr{A}(\mathbb{R}^n))^d$ can be identified by the map $\sum_{i=1}^d f_i\, e_i^* \longmapsto (f_1, \ldots, f_d)$.

Let $P_{\mathscr{A}}$ denotes the map

$$ (f_1, \ldots, f_d) \longmapsto (\sum_{i=1}^d f_i\; P_{1i}, \ldots, \sum_{i=1}^d f_i\; P_{ki}), $$

from $(\mathscr{A}(\mathbb{R}^n))^d$ to $(\mathscr{A}(\mathbb{R}^n))^k$.

Then we have the following lemma.

Lemma 3.3 *The image of* ${}^tP_{\mathscr{A}}$ *is equal to* $(\mathrm{Ker} P_{\mathscr{A}})^\circ$*, the orthogonal space of* $\mathrm{Ker} P_{\mathscr{A}}$*.*

Proof

The inclusion $\mathrm{Im}({}^tP_{\mathscr{A}}) \subset (\mathrm{Ker} P_{\mathscr{A}})^\circ$ is clear. When $\mathscr{A}(\mathbb{R}^n, E^*)$ is a Fréchet space it suffices to prove that the image of $P_{\mathscr{A}}$ is closed. Indeed, this implies that the image of $P_{\mathscr{A}}$ is a Fréchet space. Then Theorem 37.2 of [T] insures that $\mathrm{Im}({}^tP_{\mathscr{A}})$ is closed in $\mathscr{A}'(\mathbb{R}^n, E)$. On the other hand, Proposition 35.4 of [T] implies that the closure of $\mathrm{Im}({}^tP_{\mathscr{A}})$ is equal to $(\mathrm{Ker} P_{\mathscr{A}})^\circ$.

Now we will prove that the image of $P_{\mathscr{A}}$ is closed.

The case $\mathscr{A}(\mathbb{R}^n, E^*) = \mathcal{C}^\infty(\mathbb{R}^n, E^*)$ is a consequence of Theorem 0.1.1 of [B-S].

For $\mathscr{A}(\mathbb{R}^n, E^*) = \mathscr{S}(\mathbb{R}^n, E^*)$, we consider the map $\widehat{P} : (\mathcal{C}^\infty_{\overline{B}(0,1)}(\mathbb{R}^n))^d \longrightarrow (\mathcal{C}^\infty_{\overline{B}(0,1)}(\mathbb{R}^n))^k$ defined by

$$\widehat{P} = I_k \circ P_{\mathscr{S}} \circ J_d \ ,$$

where I_k is the topological isomorphism from $(\mathscr{S}(\mathbb{R}^n))^k$ to $(\mathcal{C}^\infty_{\overline{B}(0,1)}(\mathbb{R}^n))^k$ defined by :

$$I_k : (f_1, \ldots, f_k) \longmapsto (I(f_1), \ldots, I(f_k)),$$

and J_d the topological isomorphism from $(\mathcal{C}^\infty_{\overline{B}(0,1)}(\mathbb{R}^n))^d$ to $(\mathscr{S}(\mathbb{R}^n))^d$ defined by :

$$J_d : (f_1, \ldots, f_d) \longmapsto (I^{-1}(f_1), \ldots, I^{-1}(f_d)).$$

Then it is sufficient to prove that the image of $\widehat{P}$ is closed. For $f = (f_1, \ldots, f_d) \in (\mathcal{C}^\infty_{\overline{B}(0,1)}(\mathbb{R}^n))^d$, we have

$$\widehat{P}((f_1, \ldots, f_d)) = (\widehat{P}_1(f), \ldots, \widehat{P}_k(f)),$$

where, for all $1 \leq j \leq k$,

$$\widehat{P}_j(f)(x) = \begin{cases} \displaystyle\sum_{i=1}^{d} f_i(x) \ \frac{P_{ji}(x)}{(1 - ||x||^2)^{\frac{s_j}{2}}} & \text{if } x \in B(0,1) \\ 0 & \text{otherwise} \end{cases} .$$

Let $B : (\mathcal{C}^\infty_{\overline{B}(0,1)}(\mathbb{R}^n))^d \longrightarrow (\mathcal{C}^\infty_{\overline{B}(0,1)}(\mathbb{R}^n))^k$ defined by

$$B((g_1, \ldots, g_d)) = (B_1(g), \ldots, B_k(g)),$$

where

$$B_j(g)(x) = \sum_{i=1}^{d} g_i(x) \ P_{ji}(x), \qquad \text{for all } 1 \leq j \leq k.$$

Let $A : (\mathcal{C}^\infty_{\overline{B}(0,1)}(\mathbb{R}^n))^k \longrightarrow (\mathcal{C}^\infty_{\overline{B}(0,1)}(\mathbb{R}^n))^k$ defined by

$$A((g_1, \ldots, g_k)) = (A_1(g), \ldots, A_k(g)),$$

where

$$A_j(g)(x) = \begin{cases} \dfrac{g_j(x)}{(1 - ||x||^2)^{\frac{s_j}{2}}} & \text{if } x \in B(0,1) \\ 0 & \text{otherwise} \end{cases} , 1 \leq j \leq k.$$

It is clear that A is a topological isomorphism, and that $\widehat{P} = A \circ B$. Again, by Theorem 0.1.1 of [B-S], the image of the map B is closed. Consequently, the image of $\widehat{P}$ is closed.

Now we will prove the result for $\mathscr{A}(\mathbb{R}^n, E^*) = \mathcal{C}_c^\infty(\mathbb{R}^n, E^*)$ which is not a Fréchet space. Let $m \in \mathbb{N}^*$. We consider the map

$$P_m : f \longmapsto (\langle f, P_1 \rangle, \ldots, \langle f, P_k \rangle),$$

from $\mathcal{C}^\infty_{\overline{B}(0,m)}(\mathbb{R}^n, E^*)$ to $(\mathcal{C}^\infty_{\overline{B}(0,m)}(\mathbb{R}^n))^k$. We will prove that $(\mathrm{Ker} P_{\mathscr{A}})^\circ \subset \mathrm{Im}({}^t P_{\mathscr{A}})$. Let T be in $(\mathrm{Ker} P_{\mathscr{A}})^\circ$. It is clear that the restriction of T to the space $\mathcal{C}^\infty_{\overline{B}(0,m)}(\mathbb{R}^n, E^*)$ is in $(\mathrm{Ker} P_m)^\circ$. Since $\mathcal{C}^\infty_{\overline{B}(0,m)}(\mathbb{R}^n, E^*)$ is a Fréchet space, it follows, as in the first part of the proof, that $\mathrm{Im}({}^t P_m) = (\mathrm{Ker} P_m)^\circ$. So there exists a continuous linear form θ_m on $(\mathcal{C}^\infty_{\overline{B}(0,m)}(\mathbb{R}^n))^k$ such that T is equal to ${}^t P_m(\theta_m)$ on $\mathcal{C}^\infty_{\overline{B}(0,m)}(\mathbb{R}^n, E^*)$. By the Hahn-Banach theorem we extend θ_m to a continuous linear form on $(\mathcal{C}_c^\infty(\mathbb{R}^n))^k$ denoted also by θ_m. For all $m \in \mathbb{N}^*$ let

$$C_m = \{x \in \mathbb{R}^n, \ m - 2 < ||x|| < m - \frac{1}{2}\}.$$

We fix a partition of unity $(\chi_m)_{m \in \mathbb{N}^*}$ subordinated to the covering $(C_m)_{m \in \mathbb{N}^*}$ of $\mathbb{R}^n$ with $\chi_m \in \mathcal{C}_c^\infty(\mathbb{R}^n)$, and we consider the distribution $\theta = \sum_{m=1}^\infty \chi_m \, \theta_m$.

Let $f \in \mathcal{C}_c^\infty(\mathbb{R}^n, E^*)$. We have

$$\langle \theta \circ P_{\mathscr{A}}, f \rangle = \sum_{m \in \mathbb{N}^*} \langle \theta_m, \chi_m \, P_{\mathscr{A}}(f) \rangle = \sum_{m \in \mathbb{N}^*} \langle \theta_m, P_{\mathscr{A}}(\chi_m \, f) \rangle.$$

Since $\mathrm{supp}\, \chi_m \subset \overline{B}(0, m)$, then

$$\langle \theta \circ P_{\mathscr{A}}, f \rangle = \sum_{m \in \mathbb{N}^*} \langle \theta_m, P_m(\chi_m \, f) \rangle \ \text{ and } \ \langle \theta_m, P_m(\chi_m \, f) \rangle = \langle T, \chi_m \, f \rangle.$$

Consequently,

$$\langle \theta \circ P_{\mathscr{A}}, f \rangle = \langle T, \sum_{m \in \mathbb{N}^*} \chi_m \, f \rangle = \langle T, f \rangle.$$

So $T = {}^t P_{\mathscr{A}}(\theta)$. □

Lemma 3.4 *The space* $\mathscr{A}'(\mathbb{R}^n, E)^G$ *is contained in* ${}^t P_{\mathscr{A}}((\mathscr{A}(\mathbb{R}^n)^k)')$.

Proof

By Lemma 3.2 we have $\mathscr{A}'(\mathbb{R}^n, E)^G = \overline{\mathcal{C}_c^\infty(\mathbb{R}^n, E)^G}$ and by Lemma 3.3, $(\mathrm{Ker} P_{\mathscr{A}})^\circ = {}^t P_{\mathscr{A}}((\mathscr{A}(\mathbb{R}^n)^k)')$. To conclude, we need to prove the inclusion $\mathcal{C}_c^\infty(\mathbb{R}^n, E)^G \subset (\mathrm{Ker} P_{\mathscr{A}})^\circ$.

Let $\varphi \in \mathcal{C}_c^\infty(\mathbb{R}^n, E)^G$ and $f \in \mathrm{Ker} P_{\mathscr{A}}$. Then $\langle f(x), P_i(x)\rangle = 0$ for all $x \in \mathbb{R}^n$ and all $1 \leq i \leq k$. By Theorem 2.3, there exists a family of functions $\varphi_1, \dots, \varphi_k \in \mathcal{C}_c^\infty(\mathbb{R}^n)^G$ such that

$$\varphi = \sum_{i=1}^k \varphi_i \; P_i.$$

Then,

$$\langle T_\varphi, f\rangle = \int_{\mathbb{R}^n} \langle f(x), \sum_{i=1}^k \varphi_i(x) \; P_i(x)\rangle \; dx = 0.$$

□

Proof of Theorem 3.1

Let θ be in $\mathscr{A}'(\mathbb{R}^n, E)^G$. By Lemma 3.4, there exists $T \in (\mathscr{A}(\mathbb{R}^n)^k)'$ such that ${}^tP_{\mathscr{A}}(T) = \theta$. Then there is a family of distributions $(T_i)_{1\leq i\leq k}$ in $\mathscr{A}'(\mathbb{R}^n)$ such that

$$\langle T, (f_1, \dots, f_k)\rangle = \sum_{i=1}^k \langle T_i, f_i\rangle, \qquad \text{for all } f_1, \dots, f_k \in \mathscr{A}(\mathbb{R}^n),$$

and $\theta = \sum_{i=1}^k T_i \; P_i$.

For $1 \leq i \leq k$, we define, as for the G-covariant distributions in the proof of Lemma 3.2, the distribution θ_i in $\mathscr{A}'(\mathbb{R}^n)^G$ by

$$\langle \theta_i, \varphi\rangle = \int_G \langle T_i, \varphi \circ g\rangle \; d\mu(g), \qquad \text{for all } \varphi \in \mathscr{A}(\mathbb{R}^n). \tag{3.6}$$

Let f be in $\mathscr{A}(\mathbb{R}^n, E^*)$. Since θ is G-covariant, then

$$\langle \theta, f\rangle = \int_G \langle \theta, g \cdot f\rangle \; d\mu(g).$$

So we have

$$\langle \theta, f\rangle = \sum_{i=1}^k \int_G \langle T_i, \langle g \cdot f, P_i\rangle\rangle \; d\mu(g).$$

The polynomials $(P_i)_{1\leq i\leq k}$ are G-covariant, then

$$\langle T_i, \langle g \cdot f, P_i\rangle = \langle T_i, \langle f \circ g, P_i \circ g\rangle\rangle.$$

This implies by Formula (3.6) that

$$\langle \theta, f\rangle = \sum_{i=1}^k \langle \theta_i, \langle f, P_i\rangle\rangle = \langle \sum_{i=1}^k \theta_i \; P_i, f\rangle.$$

So $\theta = \sum_{i=1}^k \theta_i \; P_i$. □

References

[B] Bouaziz, A. : Intégrales orbitales sur les algèbres de Lie réductives. Invent. Math. 115, 163-207 (1994).

[BO] Bourbaki, N. : Intégration, chapitre 6. Berlin-Heidelberg : N. Bourbaki et Springer-Verlag 2007.

[B-S] Bierstone E., Schwarz G.W. : Continuous linear ivision and extension of $\mathcal{C}^\infty$ functions. Duke Math. J. 50, 233-271 (1983).

[O] Oksak, A.I. : On invariant and covariant Schwartz distributions in the case of a compact linear group. Commun. math. Phys. 14, 271-304 (1969).

[P] Poénaru, V. : Singularités $\mathcal{C}^\infty$ en présence de symétrie. Lecture Notes in Mathematics vol. 510. Berlin-Heidelberg-New York : Springer-Verlag, 1976.

[Sc] Schwartz, L. : Sur le théorème du graphe fermé. C.R. Acad. Sc. Paris. t.263, 602-605 (1966).

[T] Trèves, F. : Topological vector spaces, distributions and kernels. New york-London : Academic Press, 1967.